

A Simple Universal Generator for Continuous and Discrete Univariate T-concave Distributions

Leydold, Josef

DOI:
[10.57938/1e4101f4-abdb-4601-bd41-f5aa343ac717](https://doi.org/10.57938/1e4101f4-abdb-4601-bd41-f5aa343ac717)

Published: 01/01/2000

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):
Leydold, J. (2000). *A Simple Universal Generator for Continuous and Discrete Univariate T-concave Distributions*. (February 2000 ed.) Department of Statistics and Mathematics, Abt. f. Angewandte Statistik u. Datenverarbeitung, WU Vienna University of Economics and Business. Preprint Series / Department of Applied Statistics and Data Processing No. 30 <https://doi.org/10.57938/1e4101f4-abdb-4601-bd41-f5aa343ac717>

A Simple Universal Generator for Continuous and Discrete Univariate T-concave Distributions


Josef Leydold

Department of Applied Statistics and Data Processing
Wirtschaftsuniversität Wien

Preprint Series

Preprint 30
February 2000

<http://statmath.wu-wien.ac.at/>



A Simple Universal Generator for Continuous and Discrete Univariate T-concave Distributions

Josef Leydold

University of Economics and Business Administration, Department for Applied Statistics and Data Processing

We use inequalities to design short universal algorithms that can be used to generate random variates from large classes of univariate continuous or discrete distributions (including all log-concave distributions). The expected time is uniformly bounded over all these distributions. The algorithms can be implemented in a few lines of high level language code. In opposition to other black-box algorithms hardly any setup step is required and thus it is superior in the changing parameter case.

Categories and Subject Descriptors: G.3 [**Probability and Statistics**]: Random number generation

General Terms: Algorithms

Additional Key Words and Phrases: non-uniform random variates, universal method, ratio-of-uniforms method, transformed density rejection, discrete distributions, continuous distributions, log-concave distributions, T-concave distributions

1. INTRODUCTION

In the last decade several approaches have been introduced for so called universal (or black box) methods for generating non-uniform random variates. Recent papers propose methods where a hat function that approximates the respective probability density function or probability vector is constructed (see e.g. Ahrens [1993], Hörmann [1995], Ahrens [1995], Evans and Swartz [1998], Leydold [2000a], Leydold [2000b]; Hörmann and Derflinger [1996], Hörmann and Derflinger [1997]). These methods have (extremely) fast marginal generation time, but require a setup step, which is expensive compared to the average cost of generating one random variate. Although this setup step can be made short at the price of a much higher marginal generation time (e.g. Gilks and Wild 1992) the resulting algorithms are rather complex.

If only a few random variates are required methods like adaptive rejection sampling by Gilks and Wild [1992] or rejection from adjusted table-mountain-shaped hat functions (e.g. Hörmann 1995) have been suggested. However both require a rather expensive setup step and/or adaptation steps. Thus the approach by Devroye [1984] and Devroye [1987] is much more appropriate. It uses inequalities that holds for every log-concave distribution. It is based on the following theorems.

This work was supported by the Austrian Science Foundation (FWF), project no. P12805-MAT
Address: Augasse 2-6, A-1090 Vienna, Austria. email: Josef.Leydold@statistik.wu-wien.ac.at

THEOREM 1 (DEVROYE [1986], §VII.2.5, THEOREM 2.4). *If f is a log-concave density with mode $\mu = 0$ and $f(0) = 1$, then writing q for $F(0)$, where F denotes the c.d.f. of the distribution, we have*

$$f(x) \leq \begin{cases} \min(1, e^{1-x/(1-q)}), & (x \geq 0) \\ \min(1, e^{1+x/q}), & (x < 0) \end{cases} \quad (1)$$

The area under the bounding curve in (1) is 2.

Remark 1. If $F(\mu)$ is not known, a modified universal hat exists with area 4 (see Devroye [1986], §VII.2.3). In both cases these universal hats are not optimal. Devroye [1984] derives the properties of the optimal hat and provides a (rather expensive) generator for the corresponding density. The areas below the optimal bounding curves are $\pi^2/6$ and $\pi^2/3$, respectively, i.e., about 18% better.

THEOREM 2 (DEVROYE 1987). *For any discrete log-concave distribution with a mode at μ and probabilities p_k , we have*

$$p_{\mu+k} \leq p_\mu \min(1, e^{1-p_\mu|k|}), \quad \text{for all } k. \quad (2)$$

Remark 2. The expected number of iterations for a generator that utilizes equation (2) is $4 + p_\mu$. Devroye [1987] also gives some hints how this number can be decreased to $2 + p_\mu$ in special cases.

Algorithms that utilize these two theorems can be found in Devroye [1984] and Devroye [1987], respectively.

In this paper we introduce a new approach for universal bounding curves based on the ratio-of-uniforms method. The new algorithm is even simpler and can be applied to a larger class of distributions, including all log-concave distributions. As for Devroye's algorithm the expected number of uniform random numbers does not depend on the particular distribution. In opposition to other black-box algorithms hardly any setup step is required. Thus it is superior in the changing parameter chase.

2. CONTINUOUS DISTRIBUTIONS

2.1 Ratio-of-uniforms

The ratio-of-uniforms method introduced by Kinderman and Monahan [1977] is a flexible method that can be adjusted to a large variety of distributions. It has become a popular transformation method to generate non-uniform random variates, since it results in exact, efficient, fast and easy to implement algorithms. It is based on the following (slightly modified) theorem.

THEOREM 3 (KINDERMAN AND MONAHAN 1977). *Let $f(x)$ be a positive integrable function with support (x_0, x_1) not necessarily finite. If (V, U) is uniformly distributed in*

$$\mathcal{A} = \mathcal{A}(f) = \{(v, u): 0 < u \leq \sqrt{f(v/u + \mu)}, x_0 < v/u + \mu < x_1\}, \quad (3)$$

then $X = V/U + \mu$ has probability density function $f(x)/\int f$.

For sampling random points uniformly distributed in \mathcal{A} , rejection from a convenient enveloping region is used. Kinderman and Monahan [1977] and others use

rejection from the *minimal bounding rectangle*, i.e., the smallest possible rectangle that contains \mathcal{A} . It is given by (see Wakefield, Gelfand, and Smith [1991])

$$\mathcal{R}_{\text{mbr}} = \{(v, u): v^- \leq v \leq v^+, 0 \leq u \leq u^+\} \quad (4)$$

where

$$\begin{aligned} u^+ &= \sup_{x_0 < x < x_1} \sqrt{f(x)} \\ v^- &= \inf_{x_0 < x \leq \mu} (x - \mu) \sqrt{f(x)} \\ v^+ &= \sup_{\mu \leq x < x_1} (x - \mu) \sqrt{f(x)} \end{aligned} \quad (5)$$

2.2 A universal envelope

Assume \mathcal{A} is convex. Then it is easy to construct a universal bounding rectangle without computing these boundaries. Let μ be the mode of f , then $u^+ = \sqrt{f(\mu)}$ and \mathcal{A} has the extremal points $(0, u^+)$, (v^-, u_l) and (v^+, u_r) , for respective u_l and u_r . Define $\mathcal{A}^+ = \{(v, u) \in \mathcal{A}: v > 0\}$ and analogously \mathcal{A}^- . Then by the convexity of \mathcal{A} , the triangle with vertices at $(0, 0)$, $(0, u^+)$ and (v^+, u_r) is contained in \mathcal{A}^+ and thus has a smaller area (see figure 1). Consequently




Fig. 1. \mathcal{A} and universal bounding rectangle \mathcal{R} and squeeze \mathcal{S} for gamma(3) distribution.

$$\frac{1}{2} u^+ v^+ \leq |\mathcal{A}^+| \quad \text{and} \quad \frac{1}{2} u^+ (-v^-) \leq |\mathcal{A}^-|$$

where $|\mathcal{A}^+|$ denotes the area of \mathcal{A}^+ . From the proof of theorem 3 (see e.g. Kinderman and Monahan [1977]) it follows immediately that

$$|\mathcal{A}^-| = \frac{1}{2} \int_{x_0}^{\mu} f(x) dx = \frac{1}{2} F(\mu) \int f$$

and

$$|\mathcal{A}^+| = \frac{1}{2} \int_{\mu}^{x_1} f(x) dx = \frac{1}{2} (1 - F(\mu)) \int f.$$

Hence if the cumulate distribution function $F(x)$ at mode μ is known, we find

$$v^+ \leq (1 - F(\mu)) (ff)/\sqrt{f(\mu)} \quad \text{and} \quad v^- \geq -F(\mu) (ff)/\sqrt{f(\mu)}.$$

Therefore we can summarize our result in the following theorem.

THEOREM 4. *Suppose $\mathcal{A} = \mathcal{A}(f)$ is convex for a density function $f(x)$ with mode μ . Let F denote the c.d.f. of the distribution and let*

$$\begin{aligned} \mathcal{R} &= \{(v, u): v_l \leq v \leq v_r, 0 \leq u \leq u_m\}, \\ \mathcal{Q} &= \{(v, u): -v_m \leq v \leq v_m, 0 \leq u \leq u_m\}, \end{aligned} \quad (6)$$

where

$$\begin{aligned} u_m &= \sqrt{ff(\mu)} & v_l &= -F(\mu) v_m \\ v_m &= (ff)/\sqrt{f(\mu)} & v_r &= (1 - F(\mu)) v_m \end{aligned} \quad (7)$$

Then $\mathcal{A} \subset \mathcal{R} \subset \mathcal{Q}$ and

$$|\mathcal{R}| = 2|\mathcal{A}| \quad \text{and} \quad |\mathcal{Q}| = 4|\mathcal{A}|. \quad (8)$$

Remark 3. For the class of all distributions with convex sets \mathcal{A} , \mathcal{R} is optimal, i.e., any other universal enveloping region must contain \mathcal{R} . Analogously for \mathcal{Q} when $F(\mu)$ is not known.

Applying theorem 4 results in the following universal algorithm for distributions with convex set \mathcal{A} . It works with any multiple of the probability density function.

Algorithm SROUC

Require: p.d.f. $f(x)$, area ff , mode μ ; c.d.f. at mode $F(\mu)$ (optional)

```

/* Setup */
1:  $u_m \leftarrow \sqrt{ff(\mu)}$ ,  $v_m \leftarrow (ff)/u_m$ .
2: if  $F(\mu)$  is provided then
3:    $v_l \leftarrow -F(\mu) v_m$ ,  $v_r \leftarrow v_m + v_l$ .
4: else
5:    $v_l \leftarrow -v_m$ ,  $v_r \leftarrow v_m$ .
/* Generator */
6: repeat
7:   Generate  $U$  uniformly on  $(0, u_m)$ .
8:   Generate  $V$  uniformly on  $(v_l, v_r)$ .
9:    $X \leftarrow V/U + \mu$ .
10: until  $U^2 \leq f(X)$ .
11: return  $X$ .
```

Remark 4. By equation (8) the rejection constant of algorithm SROUC is 2 when $F(\mu)$ is known and 4 otherwise.

Remark 5. If only upper and lower bounds for ff , $F(\mu)$, $f(\mu)$ or μ are available, an accordingly modified version of algorithm SROUC still works.

2.3 A universal squeeze

When $F(\mu)$ is known we can also construct a universal squeeze.

THEOREM 5. *Suppose $\mathcal{A}(f)$ is convex for a density function $f(x)$ with mode μ . If $F(\mu)$ is given, where F denotes the c.d.f. of the distribution, then there exists a set $\mathcal{S} = \mathcal{S}(f)$, such that $\mathcal{S} \subset \mathcal{A}$. We have $(V, U) \in \mathcal{S}$ if and only if either*

$$0 \leq \frac{V}{U} \leq \frac{v_r}{u_m} \quad \text{and} \quad U v_r + V u_m \leq v_r u_m \quad (9)$$

or

$$0 \geq \frac{V}{U} \geq \frac{v_l}{u_m} \quad \text{and} \quad U v_l + V u_m \geq v_l u_m. \quad (10)$$

where u_m , v_l and v_r are as defined in theorem 4. Moreover

$$|\mathcal{S}| = |\mathcal{A}|/2. \quad (11)$$

PROOF. Let \mathcal{S} denote the universal squeeze region and assume that $\mathcal{A}^+ \neq \emptyset$. Let Δ be the triangle defined by the inequalities $v > 0$, $v/u \leq v_r/u_m$ and $u v_r + v u_m \leq v_r u_m$. Hence its vertices are $(0, 0)$, $(0, u_m)$ and $(v_r/2, u_m/2)$ (see figure 1). Define $\mathcal{S}^+ = \{(v, u) \in \mathcal{S}: v > 0\}$ and $\mathcal{R}^+ = \{(v, u) \in \mathcal{R}: v > 0\}$. Every straight line through a point $(V, U) \in \mathcal{R}^+ \setminus \mathcal{A}^+$ that does not intersect \mathcal{A}^+ , splits \mathcal{R}^+ into two parts such that (i) \mathcal{A}^+ and the edge $(0, 0)(0, u_m)$ are completely contained in the left hand part, and (ii) the area of left hand part is at least $|\mathcal{A}^+|$ and hence cannot be smaller than $|\mathcal{R}^+|/2$ (analogously to equation (8)). \mathcal{S}^+ is then the intersection of the left hand parts of all such lines. Consequently \mathcal{S}^+ must be contained in the triangles with respective vertices at $(0, 0)$, (u, u_m) and $(v_r, 0)$, and $(0, 0)$, (u, u_m) and (v_r, u_m) . Since the intersection of these triangles is given by Δ , we find $\Delta \supseteq \mathcal{S}^+$. Now notice that \mathcal{S}^+ is convex. Furthermore for every such straight line that intersects the boundary of \mathcal{R}^+ in the points $(a, 0)$ and (b, u_m) we must have $(a + b) \geq v_r/2$, since otherwise (ii) would be violated. Hence $(v_r/2, u_m/2) \in \mathcal{S}^+$, thus $\Delta \subseteq \mathcal{S}^+$ and equation (9) follows. Analogously we find \mathcal{S}^- and inequality (10) for the left hand rectangle \mathcal{R}^- . Obviously $|\mathcal{S}| = |\mathcal{S}^-| + |\mathcal{S}^+| = |\mathcal{R}|/4 = |\mathcal{A}|/2$. \square

Remark 6. For the class of all distributions with convex sets \mathcal{A} , \mathcal{S} is optimal, i.e., any other universal squeeze region is contained in \mathcal{S} .

Algorithm SR0UC can be easily extended to make use of theorem 5.

2.4 T-concave distributions

Stadlober [1989b] and Dieter [1989] have clarified the relationship of the ratio-of-uniforms method to the ordinary acceptance/rejection method. It can be viewed as rejection from a table-mountain shaped density (see figure 2). Leydold [2000a] has shown a deeper connection to the so called transformed density rejection method (see Hörmann [1995] for a description of this method). Moreover a full characterization of all distributions with convex region \mathcal{A} is derived.

THEOREM 6 (LEYDOLD 2000A). *$\mathcal{A}(f)$ is convex if and only if $f(x)$ is T-concave with transformation $T(x) = -1/\sqrt{x}$, i.e., if and only if $-1/\sqrt{f(x)}$ is a concave function.*

Notice that this class of T-concave distributions includes all log-concave distributions [Hörmann 1995].

We further can use this connection to derive universal upper and lower bounds for T-concave distributions.




Fig. 2. Universal hat and squeeze for gamma(3) distribution. Original scale (l.h.s.) and transformed scale (r.h.s.)

THEOREM 7. For any T -concave density f with $T(x) = -1/\sqrt{x}$ and mode μ let u_m, v_m, v_l and v_r be defined as in theorem 4 and let $x_m = v_m/u_m$, $x_l = v_l/u_m$ and $x_r = v_r/u_m$. Define

$$\tilde{h}(x) = \begin{cases} f(\mu) & \text{for } -x_m \leq x - \mu \leq x_m \\ \frac{v_m^2}{(x-\mu)^2} & \text{otherwise} \end{cases} \quad (12)$$

$$h(x) = \begin{cases} \frac{v_l^2}{(x-\mu)^2} & \text{for } x - \mu < x_l \\ f(\mu) & \text{for } x_l \leq x - \mu \leq x_r \\ \frac{v_r^2}{(x-\mu)^2} & \text{for } x - \mu > x_r \end{cases} \quad (13)$$

$$s(x) = \begin{cases} \left(\frac{v_l u_m}{v_l + u_m (x-\mu)} \right)^2 & \text{for } x_l \leq x - \mu < 0 \\ \left(\frac{v_r u_m}{v_r + u_m (x-\mu)} \right)^2 & \text{for } 0 \leq x - \mu < x_r \\ 0 & \text{otherwise} \end{cases} \quad (14)$$

$$\tilde{s}(x) = \begin{cases} f(\mu)/4 & \text{for } x_l \leq x - \mu \leq x_r \\ 0 & \text{otherwise} \end{cases} \quad (15)$$

Then

$$\tilde{h}(x) \geq h(x) \geq f(x) \geq s(x) \geq \tilde{s}(x) \quad \text{for all } x. \quad (16)$$

PROOF. $(v, u) \mapsto (v/u + \mu, -1/u)$ maps $\mathcal{A}(f)$ one-to-one onto the region $\mathcal{T}(f) = \{(x, y): y \leq T(f(x)) = -1/\sqrt{f(x)}, x_0 < x < x_1\}$ [Leydold 2000a]. Moreover a straight line $av + bu = c$ in $\mathcal{A}(f)$ is mapped onto the line $a(x - \mu) + cy = -b$ in $\mathcal{T}(f)$, and consequently to a curve

$$y = \left(\frac{c}{b + a(x - \mu)} \right)^2 \quad (17)$$

in the original scale, since $T^{-1}(x) = -1/x^2$. (If $c = 0$, the line $av + bu = 0$ through the origin is mapped into a line parallel to the y -axis.) We then get upper bounds by the respective boundaries of \mathcal{R} and \mathcal{Q} (with $u > 0$) in theorem 4. These are

given by $v = v_l$, $u = u_m$, $v = v_r$, and $v = -v_m$, $u = u_m$, $v = v_m$, respectively. Thus the upper bounds $\tilde{h}(x)$ and $h(x)$ follow from (17). Analogously we get $s(x)$ by theorem 5 and eq. (9), (10) and (17). The last inequality follows from the fact, that $s(x_l + \mu) = s(x_r + \mu) = (u_m/2)^2 \leq s(x)$ for all $x \in [x_l + \mu, x_r + \mu]$. \square

Remark 7. The proof of theorem 3 uses the fact that $(v, u) \mapsto (v/u + \mu, u^2)$ maps $\mathcal{A}(f)$ one-to-one onto the region $\{(x, y): y \leq f(x), x_0 < x < x_1\}$, and that it has constant Jacobian 2. An immediate consequence is that

$$\frac{1}{4} \int \tilde{h}(x) dx = \frac{1}{2} \int h(x) dx = \int f(x) dx = 2 \int s(x) dx = 4 \int \tilde{s}(x) dx. \quad (18)$$

Moreover

$$\int_{\mu}^{x_r + \mu} h(x) dx = \int_{x_r + \mu}^{\infty} h(x) dx. \quad (19)$$

Completely analogously results hold for the left hand tail of $h(x)$ and for $\tilde{h}(x)$.

We can now use eq. (16) to compile a universal generator for T -concave distributions based on the acceptance/rejection technique. Algorithm STDR generates a random variate with propability propotional to the hat function by inversion. Squeezes are omitted.

Algorithm STDR

Require: p.d.f. $f(x)$, area $\int f$, mode μ ; c.d.f. at mode $F(\mu)$ (optional)

```

/* Setup */
1:  $u_m \leftarrow \sqrt{f(\mu)}$ ,  $v_m \leftarrow (\int f)/u_m$ .
2: if  $F(\mu)$  is provided then
3:    $A \leftarrow 2 \int f$ . /* Area below hat */
4:    $a_l \leftarrow F(\mu) A/2$ ,  $a_r \leftarrow A/2 + a_l$ .
5:    $v_l \leftarrow -F(\mu) v_m$ ,  $v_r \leftarrow v_m + v_l$ .
6: else
7:    $A \leftarrow 4 \int f$ .
8:    $a_l \leftarrow A/4$ ,  $a_r \leftarrow 3A/4$ .
9:    $v_l \leftarrow -v_m$ ,  $v_r \leftarrow v_m$ .
/* Generator */
10: repeat
11:   Generate  $U$  uniformly on  $(0, A)$ .
12:   if  $U < a_l$  then
13:      $X \leftarrow -v_l^2/U + \mu$ . /* Compute  $X$  by inversion */
14:      $Y \leftarrow U^2/v_l^2$ . /* Compute  $h(X)$ . */
15:   else if  $U \leq a_r$  then
16:      $X \leftarrow v_l/u_m + (U - a_l)/u_m^2 + \mu$ .
17:      $Y \leftarrow f(\mu)$ .
18:   else /*  $U > a_r$  */
19:      $X \leftarrow v_r^2/(u_m v_r - (U - a_r)) + \mu$ .
20:      $Y \leftarrow (A - U)^2/v_r^2$ .
21:   Generate  $V$  uniformly on  $(0, 1)$ .
22: until  $VY \leq f(X)$ .

```

23: **return** X .

Remark 8. Obviously algorithm STDR is more complex (and slower) than algorithm SROUC. However it has two advantages:

(1) The rejection constant can be decreased when the domain of density is given by $(x_0, x_1) \subset \mathbb{R}$. Just replace $(0, A)$ in step 11 by (A_l, A_r) , where $A_l = \int_{-\infty}^{x_0} h(x) dx$ and $A_r = \int_{x_1}^{\infty} h(x) dx$.

(2) STDR does not suffer from the same (possible) defects when using linear congruential generators (with bad lattice structure) as have been reported for the ratio-of-uniform methods [Hörmann 1994a; Hörmann 1994b]. (However, this does not guarantee the absence of other deficiencies.)

2.5 The mirror principle

Devroye [1984] suggests the usage of a hat function for $f(x) + f(-x)$ when $F(\mu)$ is not known to reduce the expected number of uniform random numbers. To apply this idea to our situation we need the following result.

LEMMA 8. *Let $g_1(x)$ and $g_2(x)$ be two non-negative functions with respective bounding rectangles \mathcal{R}_1 and \mathcal{R}_2 for $\mathcal{A}(g_1)$ and $\mathcal{A}(g_2)$ with common left lower vertex $(0, 0)$ and the respective right upper vertices (v_1, u_1) and (v_2, u_2) . Then*

$$\mathcal{A}(g_1 + g_2) \subseteq \left\{ (v, u) : 0 \leq v \leq \sqrt{v_1^2 + v_2^2}, 0 < u \leq \sqrt{u_1^2 + u_2^2} \right\}.$$

PROOF. Let $(v, u) \in \mathcal{A}(g_1 + g_2)$. Obviously $v \geq 0$. By equation (5) $u^2 \leq \sup(g_1(x) + g_2(x)) \leq \sup g_1(x) + \sup g_2(x) \leq u_1^2 + u_2^2$ and $v^2 \leq \sup(x - \mu)^2 (g_1(x) + g_2(x)) \leq \sup(x - \mu)^2 g_1(x) + \sup(x - \mu)^2 g_2(x) \leq v_1^2 + v_2^2$ as proposed. \square

THEOREM 9. *For any T -concave density f with $T(x) = -1/\sqrt{x}$ and mode μ let*

$$\widehat{\mathcal{R}} = \{(v, u) : -v_m \leq v \leq v_m, 0 < u \leq \sqrt{2} u_m\} \quad (20)$$

where v_m and u_m are as defined in theorem 4. Then

$$\mathcal{A}(f(x) + f(-x)) \subset \widehat{\mathcal{R}} \quad \text{and} \quad |\widehat{\mathcal{R}}| = 4\sqrt{2} |\mathcal{A}(f)|. \quad (21)$$

PROOF. Let $\mathcal{R}(p) = \{(v, u) : -pv_m \leq v \leq (1-p)v_m, 0 < u \leq u_m\}$. Then by theorem 4 we find $\mathcal{A}(f(x)) \subset \mathcal{R}(F(\mu))$ and $\mathcal{A}(f(-x)) \subset \mathcal{R}(1 - F(\mu))$. Hence by lemma 8 $\mathcal{A}(f(x) + f(-x)) \subset \bigcup_{p \in [0,1]} \{(v, u) : -v(p) \leq v \leq v(p), 0 < u \leq \sqrt{2} u_m\} = \widehat{\mathcal{R}}$, where $v(p) = \sqrt{p^2 v_m^2 + (1-p)^2 v_m^2}$. Equation (21) follows immediately from equation (8). \square

Remark 9. $\widehat{\mathcal{R}}$ is not optimal. However the optimal envelope $\widehat{\mathcal{R}}_{\text{opt}}$ is not rectangular and contains the rectangles $\{(v, u) : -v_m/2 \leq v \leq v_m/2, 0 < u \leq \sqrt{2} u_m\}$ and $\{(v, u) : -v_m \leq v \leq v_m, 0 < u \leq u_m\}$. Thus we find the estimate $|\widehat{\mathcal{R}}_{\text{opt}}| \geq (1 - \frac{1}{2}(\sqrt{2} - 1)) |\widehat{\mathcal{R}}| \geq 0.79 |\widehat{\mathcal{R}}|$.

Using theorem 9 we can compile the following algorithm. It reduces the expected number of uniform random variates at the expense of more evaluations of $f(x)$.

Algorithm SROUCM

Require: p.d.f. $f(x)$, area $\int f$, mode μ .

```

/* Setup */
1:  $u_m \leftarrow \sqrt{f(\mu)}$ ,  $v_m \leftarrow (\int f)/u_m$ .
/* Generator */
2: loop
3:   Generate  $U$  uniformly on  $(0, \sqrt{2} u_m)$ .
4:   Generate  $V$  uniformly on  $(-v_m, v_m)$ .
5:    $X \leftarrow V/U$ .
6:   if  $U^2 \leq f(X + \mu)$  then
7:     return  $X + \mu$ .
8:   if  $U^2 \leq f(X + \mu) + f(-X + \mu)$  then
9:     return  $-X + \mu$ .

```

Remark 10. By equation (21) the rejection constant of algorithm SROUCM is $2\sqrt{2}$ in opposition to 4 in algorithm SROUC when $F(\mu)$ is not known.

3. DISCRETE DISTRIBUTIONS

Stadlober [1989a] has shown that the ratio-of-uniforms method is well suited for generating from discrete distributions. Indeed, considerations from §2 can also be used to design a universal algorithm for discrete distributions. However some modifications are necessary.

A discrete distribution with probability vector p_i , with support $I \subseteq \mathbb{Z}$, is called *T-concave* if

$$p_i \geq \frac{1}{2}(T(p_{i-1}) + T(p_{i+1})) \quad \text{for all } i \in I \tag{22}$$

For log-concave distributions we have $T(x) = \log(x)$ and $p_i^2 \geq p_{i-1} p_{i+1}$. Obviously p_i is unimodal. Denote its mode by μ . For the following assume that p_i is *T-concave* with transformation $T(x) = -1/\sqrt{x}$. Let

$$f_p(x) = \begin{cases} p_{\lfloor x \rfloor} & \text{for } \lfloor x \rfloor \in I \\ 0 & \text{otherwise} \end{cases} \tag{23}$$

where $\lfloor x \rfloor$ denotes the largest integer not greater than x . Since f_p is a step function, $\mathcal{A}(f_p)$ cannot be convex. Consider the convex hull \mathcal{C} of \mathcal{A} . Because of inequality (22), \mathcal{C} contains the points $((i+1-\mu)\sqrt{p_i}, \sqrt{p_i})$ for all $i \geq \mu$, and $((i-\mu)\sqrt{p_i}, \sqrt{p_i})$ for all $i \leq \mu$, with $i \in I \subseteq \mathbb{Z}$ (use transformation $(v, u) \mapsto (v/u + \mu, u^2)$, see remark after the proof of theorem 7). These are the “spikes” of \mathcal{A} . Let (v^+, u_r) be the right extremal point of \mathcal{A} (see (5)).

LEMMA 10. *Let Δ be the triangle with vertices at $(0, 0)$, $(0, u^+)$ and (v^+, u_r) . Then $|\Delta| \leq |\mathcal{A}^+|$, where $\mathcal{A}^+ = \{(v, u) \in \mathcal{A} : v > 0\}$.*

PROOF. Notice that the edges $(0, 0)(0, u^+)$ and $(0, 0)(v^+, u_r)$ are always contained in the closure of \mathcal{A} . Moreover the third edge $(0, u^+)(v^+, u_r)$ is also contained in \mathcal{A} whenever $u_r = 0$ by inequality (22). Then $\Delta \subseteq \mathcal{A}$ and the proposition follows. Now assume $u_r > 0$. Edge $(0, u^+)(v^+, u_r)$ is contained in the quadrangle \mathcal{Q} with vertices in $(0, u^+)$, (u^+, u^+) , (v^+, u_r) and $(v^+ - u_r, u_r)$. Figure 3 shows the

“worst case” where equality holds in (22) for all $i \in (\mu, \mu + v^+)$. Notice that edge $(0, u^+)(v^+, u_r)$ splits Q into two parts with $|\Delta \cap Q| \leq |Q|/2$, because $u_r \leq u^+$. Moreover Q in figure 3 can be partitioned into quadrangles each with two sides parallel to the v -axis. In each of these quadrangles the region that is contained in \mathcal{A} is larger than its complement. Hence $|\mathcal{A} \cap Q| \geq |Q|/2$ and the proposition follows. \square




Fig. 3. Quadrangle Q and edge $(0, u^+)(v^+, u_r)$

It obvious that an analogous result holds for \mathcal{A}^- and we arrive at the following proposition. Notice that $\sup_{i < \mu} p_i = p_{\mu-1}$.

THEOREM 11. *Let p_i , $i \in \mathbb{Z}$, be a T -concave probability vector of a discrete distribution, with $T(x) = -1/\sqrt{x}$ and mode μ . Let F denote the c.d.f. of the distribution and let $\mathcal{R}_d = \mathcal{R}_d^- \cup \mathcal{R}_d^+$ with*

$$\begin{aligned} \mathcal{R}_d^- &= \{(v, u): -F(\mu-1) \sum p_i / \sqrt{p_{\mu-1}} \leq v \leq 0, 0 \leq u \leq \sqrt{p_{\mu-1}}\} \\ \mathcal{R}_d^+ &= \{(v, u): 0 \leq v \leq (1 - F(\mu)) \sum p_i / \sqrt{p_\mu}, 0 \leq u \leq \sqrt{p_\mu}\} \end{aligned} \quad (24)$$

and $\mathcal{Q}_d = \mathcal{Q}_d^- \cup \mathcal{Q}_d^+$ with

$$\begin{aligned} \mathcal{Q}_d^- &= \{(v, u): -\sum p_i / \sqrt{p_{\mu-1}} \leq v \leq 0, 0 \leq u \leq \sqrt{p_{\mu-1}}\} \\ \mathcal{Q}_d^+ &= \{(v, u): 0 \leq v \leq \sum p_i / \sqrt{p_\mu}, 0 \leq u \leq \sqrt{p_\mu}\} \end{aligned} \quad (25)$$

Then $\mathcal{A} \subset \mathcal{R}_d \subset \mathcal{Q}_d$ and

$$|\mathcal{R}_d| = 2|\mathcal{A}| \quad \text{and} \quad |\mathcal{Q}_d| = 4|\mathcal{A}|. \quad (26)$$

Remark 11. We set $\mathcal{R}_d^- = \emptyset$ whenever $p_{\mu-1} = 0$.

Algorithm SROUD

Require: p.v. p_i , area $\sum p_i$, mode μ ; c.d.f. at mode $F(\mu)$ (optional)

/* Setup */

- 1: $u_l \leftarrow \sqrt{p_{\mu-1}}$, $u_r \leftarrow \sqrt{p_\mu}$.
- 2: **if** $F(\mu)$ is provided **then**

```

3:   $v_l \leftarrow -F(\mu) \sum p_i/u_l, \quad v_r \leftarrow (1 - F(\mu - 1)) \sum p_i/u_r.$ 
4:  else
5:   $v_l \leftarrow -\sum p_i/u_l, \quad v_r \leftarrow \sum p_i/u_r.$ 
   /* Generator */
6:  repeat
7:    Generate  $V$  uniformly on  $(v_l, v_r)$ .
8:    if  $V < 0$  then
9:      Generate  $U$  uniformly on  $(0, u_l)$ .
10:   else
11:     Generate  $U$  uniformly on  $(0, u_r)$ .
12:    $I \leftarrow \lfloor V/U \rfloor + \mu.$ 
13: until  $U^2 \leq p_I.$ 
14: return  $I.$ 

```

Remark 12. The rejection constant of algorithm `SR0UD` is 2 when $F(\mu)$ is known and 4 otherwise.

Remark 13. Obviously the mirror principle can also be applied to the discrete case.

4. COMPUTATIONAL EXPERIENCES

We have coded versions of `srouc` (with and without using the universal squeeze), `sroucm` (i.e., using the mirror principle) and of `stdr` (with and without using squeeze $\tilde{s}(x)$) where we have restricted the domain of the hat to the domain of the given p.d.f. It is obvious that the expected number of uniform random numbers for new algorithms is quite high and the marginal generation time is higher than that for specialized algorithms or fast universal algorithm that require a more expensive setup step (e.g., algorithm `arou` in Leydold [2000a], or `utdr` in [Hörmann 1995], that uses a semi-empirical rule to construct a hat function). However, when only a couple of random numbers are requested, the new algorithms are superior both in generation time and the size and complexity of their codes.

To get an idea of the performance we ran several tests. The expected number of uniform random numbers for both `srouc` and `stdr` is 8 when the c.d.f. at the mode is not known and 4 otherwise. When using the mirror principle in the first case, it could be reduced to 5.66. However its usage is only recommended when the generation of uniform random numbers is (very) expensive compared to the evaluation of the p.d.f. As expected `srouc` is a little bit faster than `stdr`. However when the domain of the head function is restricted to the domain of the p.d.f. `stdr` requires less iterations and less uniform random numbers (e.g. 3.26 for the beta(5,7) distribution) and is thus faster.

When comparing the total times (including setup) for generating 10 random variates we found that both algorithms have about the same performance as `utdr` when c.d.f. at the mode is known (and thus are faster, when less random variates are required). If the c.d.f. at the mode is not known, they are at least faster than `arou` or the algorithm by Gilks and Wild [1992].

REFERENCES

- AHRENS, J. H. 1993. Sampling from general distributions by suboptimal division of domains. *Grazer Math. Berichte* 319, 20 pp.
- AHRENS, J. H. 1995. A one-table method for sampling from continuous and discrete distributions. *Computing* 54, 2, 127–146.
- DEVROYE, L. 1984. A simple algorithm for generating random variates with a log-concave density. *Computing* 33, 3–4, 247–257.
- DEVROYE, L. 1986. *Non-Uniform Random Variate Generation*. Springer-Verlag, New-York.
- DEVROYE, L. 1987. A simple generator for discrete log-concave distributions. *Computing* 39, 87–91.
- DIETER, U. 1989. Mathematical aspects of various methods for sampling from classical distributions. In E. A. MC NAIR, K. J. MUSSELMAN, AND P. HEIDELBERGER Eds., *Proc. 1989 Winter Simulation Conf.*, pp. 477–483.
- EVANS, M. AND SWARTZ, T. 1998. Random variable generation using concavity properties of transformed densities. *Journal of Computational and Graphical Statistics* 7, 4, 514–528.
- GILKS, W. R. AND WILD, P. 1992. Adaptive rejection sampling for Gibbs sampling. *Applied Statistics* 41, 2, 337–348.
- HÖRMANN, W. 1994a. A note on the quality of random variates generated by the ratio of uniforms method. *ACM TOMACS* 4, 1, 96–106.
- HÖRMANN, W. 1994b. The quality of non-uniform random numbers. In H. DYCKHOFF, U. DERINGS, M. SALOMON, AND H. C. TIJMS Eds., *Operations Research Proceedings 1993* (Berlin, 1994), pp. 329–335. Springer Verlag.
- HÖRMANN, W. 1995. A rejection technique for sampling from T-concave distributions. *ACM Trans. Math. Software* 21, 2, 182–193.
- HÖRMANN, W. AND DERFLINGER, G. 1996. Rejection-inversion to generate variates from monotone discrete distributions. *ACM TOMACS* 6, 3, 169–184.
- HÖRMANN, W. AND DERFLINGER, G. 1997. An automatic generator for a large class of unimodal discrete distributions. In A. R. KAYLAN AND A. LEHMANN Eds., *ESM 97* (1997), pp. 139–144.
- KINDERMAN, A. J. AND MONAHAN, F. J. 1977. Computer generation of random variables using the ratio of uniform deviates. *ACM Trans. Math. Software* 3, 3, 257–260.
- LEYDOLD, J. 2000a. Automatic sampling with the ratio-of-uniforms method. *ACM Trans. Math. Software*. to appear.
- LEYDOLD, J. 2000b. A note on transformed density rejection. *Computing*. to appear.
- STADLOBER, E. 1989a. Ratio of uniforms as a convenient method for sampling from classical discrete distributions. In E. A. M. ET AL. Ed., *Proc. 1989 Winter Simulation Conf.*, pp. 484–489.
- STADLOBER, E. 1989b. *Sampling from Poisson, binomial and hypergeometric distributions: Ratio of uniforms as a simple and fast alternative*. Number 303 in Bericht der Mathematisch-Statistischen Sektion in der Forschungsgesellschaft Joanneum-Graz.
- WAKEFIELD, J. C., GELFAND, A. E., AND SMITH, A. F. M. 1991. Efficient generation of random variates via the ratio-of-uniforms method. *Statist. Comput.* 1, 129–133.